



Cite this: *Chem. Commun.*, 2016, 52, 4891

Received 27th November 2015,
Accepted 28th February 2016

DOI: 10.1039/c5cc09817c

www.rsc.org/chemcomm

Copper-catalyzed cross-coupling reactions of epoxides with *gem*-diborylmethane: access to γ -hydroxyl boronic esters†

Ahmed Ebrahim-Alkhalil, Zhen-Qi Zhang, Tian-Jun Gong, Wei Su, Xiao-Yu Lu, Bin Xiao* and Yao Fu*

Herein, we describe a novel copper-catalyzed epoxide opening reaction with *gem*-diborylmethane. Aliphatic, aromatic epoxides as well as aziridines are converted to the corresponding γ -pinacolboronate alcohols or amines in moderate to excellent yields. This new reaction provides beneficial applications for classic epoxide substrates as well as interesting *gem*-diborylalkane reagents.

Transition metal catalyzed cross-coupling reactions towards the formation of C(sp³)-C(sp³) bonds play a significant role in organic synthesis,¹ whereas, in these reactions, alkyl halides and pseudo-halides were the common electrophiles.² As a class of functional electrophiles, epoxide compounds could undergo ring-opening cross-coupling and result in the corresponding β -substituted alcohols.³ Compared to traditional organometallic reagents⁴ (e.g. Grignard reagents, organozinc reagents or organolithium reagents), cross-coupling of epoxides with less reactive nucleophiles,⁵ in particular organoborons,⁶ has received increasing attention. For instance, Doyle has presented nickel-catalyzed Suzuki-type cross-coupling of epoxides with a variety of aryl boronic acids, generating rearranged products *via* a multicatalytic process.^{6b} Very recently, our research group realized the copper-catalyzed terminal direct ring-opening coupling of epoxides with aryl boronic esters (Scheme 1a).^{6c}

Regarding another research interest of our group, *gem*-diborylalkanes⁷ emerged as a kind of valuable compounds due to their ability to introduce a boron group (*i.e.*, Bpin, pinacol boronic esters),⁸ in C-C bond-forming reactions, which have more derived possibilities in further organic reactions. Herein, we report our efforts to expand the epoxide ring-opening reaction of our previously reported method to include the *gem*-diborylmethane derivative (Scheme 1b).

This study includes a series of aliphatic and aromatic epoxides which were converted to the corresponding secondary

a. Previous work



b. This work




Scheme 1 Previous studies on transition metal catalyzed Suzuki Miyaura cross-coupling reaction of epoxides.

or tertiary γ -hydroxyl boronic esters. In addition, *N*-sulfonyl aziridines were also suitable for this transformation to afford the corresponding γ -amino boronic esters. Thus, this new reaction provides valuable applications for classic epoxide substrates as well as interesting *gem*-diborylalkane reagents.

To verify our initial copper catalyst hypothesis, we started the reaction by choosing 2-(phenoxy)methyl oxirane and diborylmethane as the model substrates at 60 °C in argon. On the basis of previous studies on Cu-catalyzed cross-coupling,^{2c,6c,7g,9} various bases were subsequently screened in this reaction, such as NaO^tBu, KO^tBu, K₃PO₄, Cs₂CO₃, LiOMe and LiO^tBu (Table 1, entries 1–8), among which LiO^tBu was found to be the optimal one and afforded product **3a**, albeit in low yield (entry 9). The process occurred smoothly in 1,4-dioxane with CuI as catalyst, loading the expected product **3a** in moderate yield (entry 10). To our delight, in THF, CuI performed well with high effectivity delivering the deborylative alkylation alcohol product in high yield (88% GC yield and 85% isolated yield, entry 11). The decrease of copper catalyst loading to 10 mol% led to a slightly lower yield of 75% (entry 12). Next, a trace product was detected in the absence of CuI (entry 13). Furthermore, the use of CuCl in this transformation afforded lower yield (entry 14). Finally, considering our previously studies on using potassium iodide to promote the cross-coupling reaction,^{6c} the iodide source (potassium iodide or Bu₄NI) was also examined, the results indicated that the addition of iodide salt did not accelerate this conversion.

Hefei National Laboratory for Physical Sciences at the Microscale, iChEM, CAS Key Laboratory of Urban Pollutant Conversion, Anhui Province Key Laboratory of Biomass Clean Energy, University of Science and Technology of China, Hefei 230026, China. E-mail: binxiao@ustc.edu.cn, fuyao@ustc.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5cc09817c



Table 1 Optimization of reaction conditions^a


Entry	[Cu]	Base	Additive	Solvent	Yield ^a (%)
1	CuI	NaO ^t Bu	—	THF	Trace
2	CuI	KO ^t Bu	—	THF	n.r
3	CuI	NaO ^t Am	—	THF	Trace
4	CuI	K ₃ PO ₄	—	THF	n.r
5	CuI	Cs ₂ CO ₃	—	THF	n.r
6	CuI	LiOMe	—	THF	Trace
7	CuI	LiO ^t Bu	—	CH ₃ CN	Trace
8	CuI	LiO ^t Bu	—	Toluene	Trace
9	CuI	LiO ^t Bu	—	DMF	21
10	CuI	LiO ^t Bu	—	Dioxane	57
11 ^{b,c}	CuI	LiO^tBu	—	THF	88(85^d)
12 ^e	CuI	LiO ^t Bu	—	THF	75
13	—	LiO ^t Bu	—	THF	Trace
14	CuCl	LiO ^t Bu	—	THF	44
15 ^f	CuCl	LiO ^t Bu	Bu ₄ NI	THF	Trace
16 ^g	CuCl	LiO ^t Bu	KI	THF	75
17 ^h	CuCl	LiO ^t Bu	KI	THF	16

^a Reaction conditions: epoxide (0.2 mmol), diborylmethane (2 equiv.), CuI (20 mol%), base (3 equiv.), in 0.5 mL of solvent at 60 °C for 24 h under an Ar atmosphere. The yield was determined by GC using benzophenone as internal standard (average of two GC runs). ^b At 80 °C, GC yield was 63%. ^c Reaction was performed for 18 h, yield = 69%. ^d Isolated yield. ^e CuI (10 mol%) was used. ^f Bu₄NI (20 mol%) was added. ^g KI (20 mol%) was added. ^h KI (1.5 equiv.) was used. DMF = *N,N*-dimethylformamide.

With the optimal reaction conditions in hand, various epoxides were subjected to examination of the generality of this new method (Table 2). First, as a series of analogues of **1a**, we continued our new reaction to other protected glycidyl substrates. A variety of aryl (**3a–i**, **k**), benzyl (**3j**) and other alkyl (**3m**) substrates effectively participated in the reaction and afforded the products in good yields. The aromatic ring with the functional groups substituted at the *ortho*-, *meta*-, *para*-position did not affect the reaction results. Next, it is worth noting that the substrates containing aryl chloride and bromide can be obtained smoothly; wherein, these halides are susceptible for further transformations by metal catalysis. Furthermore, this cross-coupling reaction was not limited to glycidyl substrates, and non-glycidyl substrates (**3l**, **3o**, **3n**, **3p**) were also suitable substrates. For instance, a heteroatom substituent on the side chain, such as sulfur (**3o**), and nitrogen (**3n**) were tolerable in this reaction. Finally, we found that the more active aromatic epoxide (**3p**) was the suitable substrate under our standard reaction conditions, which was not tolerated in our previous work on Cu-catalyzed cross-coupling of epoxides with arylboronates.^{6c}

Furthermore, 1,1-disubstituted epoxides were also used in this reaction and converted to tertiary alcohol products (Table 3). Aromatic compounds with *para*-substituted (*e.g.* electron rich aromatic PMP **3ba**), naphthyl (**3bb**) and amide (**3bc**) groups were coupled well under standard conditions to give the desired products in moderate to good yields (59–74% yield). The ketal group survived in this transformation and afforded the corresponding product (**3bd**) in a yield of 60%, at a higher reaction temperature (*i.e.* 80 °C). The ring-opening cross-coupling of these compounds showed

Table 2 Scope of epoxides^{a,b}



R: H **3a**, 85%
p-OMe **3b**, 81%
m-OMe **3c**, 56%
p-Cl **3d**, 81%
o-F **3e**, 78%
p-I **3f**, 67%

3g, 73%
3h, 68%
3i, 72%
3j, 67%
3k, 79%
3l, 79%
3m, 77%
3n, 70%
3o, 81%
3p, 57%



^a Reaction conditions: epoxides (0.2 mmol), diborylmethane (2 equiv.), CuI (0.04 mmol), LiO^tBu (0.6 mmol), in 0.5 mL of THF at 60 °C for 24 h under an Ar atmosphere. ^b Isolated yield.

identical terminal-selectivity to our previous studies.^{6c} However, 1,2-disubstituted epoxides did not afford the desired product.


The generation of alkylboronic ester derivatives, in particular γ -pinacolboronate amines, is important in organic synthesis, which have numerous transformations and pharmaceutical applications.¹⁰ To our delight, under the standard reaction conditions, 2-butyl-1-tosylaziridine converted smoothly and provided the corresponding amine (**3ca**) in good yield (Scheme 2, 79% yield).

Due to the importance of chiral compounds in pharmaceutical chemistry, additional application of this reaction has been achieved for (*S*)-2-((benzyloxy)methyl)oxirane which reacted well with diborylmethane, followed by oxidation step with basic H₂O₂ at lower temperature,¹¹ to give the 1,3-diol product in high yield with fully maintained configuration as illustrated in Scheme 3.

In conclusion, we have developed the first copper-catalyzed ring opening coupling of epoxides with *gem*-diborylmethane

Table 3 Scope of 1,1-disubstituted epoxides^{a,b}

Scheme 2 Copper-catalyzed cross-coupling of aziridine derivatives.



Scheme 3 Copper-catalyzed cross-coupling of chiral epoxides.

reagents. In addition, *N*-sulfonyl aziridine could also be converted in this transformation. This newly developed reaction provided straightforward access to γ -hydroxyl boronic esters which are important synthetic intermediates in C–C bond-forming reactions. Furthermore, this strategy extended the applications of classic organic synthon epoxides as well as novel *gem*-diborylmethane reagents. The mechanism study for this reaction and transformation of more challenging substituted bis(boronates) are currently underway in our laboratory.

This work was financially supported by the 973 Program (2012CB215306), NSFC(21325208, 21361140372, 21472181, 21572212), IPDFHCPST (2014FXCX006), CAS (KFJ-EW-ST5-051, YIPA-2015371), FANEDD (201424), FRFCU, PCSIRT and CAS-TWAS President's Fellowship for International PhD Students.

Notes and references

- 1 F. Diederich and P. J. Stang, *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH, Weinheim, 1998.
- 2 (a) Z. Lu and G. C. Fu, *Angew. Chem., Int. Ed.*, 2010, **49**, 6676; (b) S. L. Zultanski and G. C. Fu, *J. Am. Chem. Soc.*, 2011, **133**, 15362; (c) C. T.

- Yang, Z. Q. Zhang, Y. C. Liu and L. Liu, *Angew. Chem., Int. Ed.*, 2011, **50**, 3904; (d) A. Wilsily, F. Tramutola, N. A. Owston and G. C. Fu, *J. Am. Chem. Soc.*, 2012, **134**, 5794; (e) T. Hatakeyama, T. Hashimoto, K. K. A. D. S. Kathiriarachchi, T. Zenmyo, H. Seike and M. Nakamura, *Angew. Chem., Int. Ed.*, 2012, **51**, 8834; (f) C.-T. Yang, Z.-Q. Zhang, J. Liang, J.-H. Liu, X.-Y. Lu, H.-H. Chen and L. Liu, *J. Am. Chem. Soc.*, 2012, **134**, 11124; (g) M. Kuriyama, M. Shinozawa, N. Hamaguchi, S. Matsuo and O. Onomura, *J. Org. Chem.*, 2014, **79**, 5921.
- (a) D. M. Hodgson and C. D. Bray, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 145–184; (b) P. Crotti and M. Pineschi, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 271–313; (c) B. Olofsson and P. Somfai, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 315–347; (d) V. V. Fokin and P. Wu, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 443–477; (e) J. A. Kalow and A. G. Doyle, *J. Am. Chem. Soc.*, 2010, **132**, 3268; (f) J. A. Kalow and A. G. Doyle, *J. Am. Chem. Soc.*, 2011, **133**, 16001; (g) J. A. Kalow, D. E. Schmitt and A. G. Doyle, *J. Org. Chem.*, 2012, **77**, 4177; (h) T. J. A. Graham, R. F. Lambert, K. Ploessl, H. F. Kung and A. G. Doyle, *J. Am. Chem. Soc.*, 2014, **136**, 5291; (i) Y. Zhao and D. J. Weix, *J. Am. Chem. Soc.*, 2014, **136**, 48; (j) C.-Y. Huang and A. G. Doyle, *Chem. Rev.*, 2014, **114**, 8153; (k) Y. Zhao and D. J. Weix, *J. Am. Chem. Soc.*, 2015, **137**, 3237.
- (a) C. Bonini, L. Chiummiento, M. T. Lopardo, M. Pullez, F. Colobert and G. Solladié, *Tetrahedron Lett.*, 2003, **44**, 2695; (b) E. Vrancken, A. Alexakis and P. Mangeney, *Eur. J. Org. Chem.*, 2005, 1354; (c) M. Pineschi, *Eur. J. Org. Chem.*, 2006, 4979; (d) M. Alam, C. Wise, C. A. Baxter, E. Cleator and A. Walkinshaw, *Org. Process Res. Dev.*, 2012, **16**, 435.
- L. P. C. Nielsen and E. N. Jacobsen, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 229–269.
- (a) J. Kjellgren, J. Aydin, O. A. Wallner, I. V. Saltanova and K. J. Szabó, *Chem. – Eur. J.*, 2005, **11**, 5260; (b) D. K. Nielsen and A. G. Doyle, *Angew. Chem., Int. Ed.*, 2011, **50**, 6056; (c) X. Lu, C. Yang, J. Liu, Z. Zhang, X. Lu, X. Lou, B. Xiao and Y. Fu, *Chem. Commun.*, 2015, **51**, 2388.
- (a) K. Endo, T. Ohkubo, M. Hirokami and T. Shibata, *J. Am. Chem. Soc.*, 2010, **132**, 11033; (b) K. Endo, M. Hirokami and T. Shibata, *J. Org. Chem.*, 2010, **75**, 3469; (c) K. Endo, T. Ohkubo and T. Shibata, *Org. Lett.*, 2011, **13**, 3368; (d) K. Endo, T. Ohkubo, T. Ishioka and T. Shibata, *J. Org. Chem.*, 2012, **77**, 4826; (e) K. Endo, T. Ishioka, T. Ohkubo and T. Shibata, *J. Org. Chem.*, 2012, **77**, 7223; (f) K. Endo and T. Ishioka, *Synlett*, 2014, 2184; (g) Z. Zhang, C. Yang, L. Liang, B. Xiao, X. Lu, J. Liu, Y. Sun, T. B. Marder and Y. Fu, *Org. Lett.*, 2014, **16**, 6342.
- (a) C. Sun, B. Potter and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 6534; (b) K. Hong, X. Liu and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 10581; (c) B. Potter, A. Szymaniak, E. Edelstein and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 17918; (d) J. R. Coombs, L. Zhang and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 16140; (e) J. R. Coombs, L. Zhang and J. P. Morken, *Org. Lett.*, 2015, **17**, 1708; (f) M. V. Joannou, B. S. Moyer and S. J. Meek, *J. Am. Chem. Soc.*, 2015, **137**, 6176.
- Y.-Y. Sun, J. Yi, X. Lu, Z.-Q. Zhang, B. Xiao and Y. Fu, *Chem. Commun.*, 2014, **50**, 11060.
- (a) M. A. Beenen, C. An and J. A. Ellman, *J. Am. Chem. Soc.*, 2008, **130**, 6910; (b) L. J. Milo, J. J. H. Lai, W. Wu, Y. Liu, H. Maw, Y. Li, Z. Jin, Y. Shu, S. E. Poplawski, Y. Wu, D. G. Sanford, J. L. Sudmeier and W. Bachovchin, *J. Med. Chem.*, 2011, **54**, 4365 and references therein; (c) A. P. Pulis and V. K. Aggarwal, *J. Am. Chem. Soc.*, 2012, **134**, 7570; (d) X. Lu, J. Yi, Z.-Q. Zhang, J.-J. Dai, J.-H. Liu, B. Xiao, Y. Fu and L. Liu, *Chem. – Eur. J.*, 2014, **20**, 15339 and references therein; (e) C. Sandford, R. Rasappan and V. K. Aggarwal, *J. Am. Chem. Soc.*, 2015, **137**, 10100; (f) J. Llaveria, D. Leonori and V. K. Aggarwal, *J. Am. Chem. Soc.*, 2015, **137**, 10958.
- (a) H. Y. Cho and J. P. Morken, *J. Am. Chem. Soc.*, 2008, **130**, 16140; (b) H. Y. Cho and J. P. Morken, *J. Am. Chem. Soc.*, 2010, **132**, 7576; (c) L. T. Kliman, S. N. Mlynarski, G. E. Ferris and J. P. Morken, *Angew. Chem., Int. Ed.*, 2012, **51**, 521; (d) C. Yang, Z. Zhang, H. Tajuddin, C. Wu, J. Liang, J. Liu, Y. Fu, M. Czystewska, P. G. Steel, T. B. Marder and L. Liu, *Angew. Chem., Int. Ed.*, 2012, **51**, 528; (e) T. P. Blaisdell, T. C. Caya, L. Zhang, A. Sanz-Marco and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 9264.